

AN ENVELOPING SERIES FOR THE ZETA FUNCTION

BY

J. N. FRANKLIN

(Communicated by Prof. J. G. VAN DER CORPUT at the meeting of Sept. 27, 1958)

The concept of an enveloping series was introduced by G. PÓLYA and G. SZEGÖ [1]. A convergent or divergent series of terms a_n is said to envelop a number A if for each $n=0, 1, \dots$

$$(1) \quad A - (a_0 + a_1 + \dots + a_n) = \theta_n a_{n+1} \text{ with } 0 < \theta_n < 1.$$

In this case we write

$$(2) \quad A \propto \sum_{n=0}^{\infty} a_n.$$

A similar notion had been developed by G. A. SCHOTT and G. N. WATSON [2]. The concept has been generalized and applied extensively by J. G. VAN DER CORPUT [3].

The purpose of this paper is to obtain an enveloping series for the function

$$(3) \quad \zeta(s) - \sum_{n=1}^{a-1} n^{-s} = \zeta(s, a),$$

where a is a positive integer and $\zeta(s)$ is the zeta function of RIEMANN. The function $\zeta(s, a)$ is the zeta function of HURWITZ [4], p. 265.

Let $\text{Re } s = \sigma$. We have [4], p. 266,

$$(4) \quad \zeta(s, a) \Gamma(s) = \int_0^{\infty} \frac{x^{s-1} e^{-ax}}{1 - e^{-x}} dx \quad (\sigma > 1).$$

Using the identity

$$(5) \quad (1 - e^{-x})^{-1} = \frac{1}{x} + \frac{1}{2} + \left(-\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} \right),$$

we find

$$(6) \quad \zeta(s, a) \Gamma(s) = \frac{\Gamma(s-1)}{a^{s-1}} + \frac{1}{2} \frac{\Gamma(s)}{a^s} + \int_0^{\infty} e^{-ax} x^{s-1} \left(-\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} \right) dx.$$

By analytic continuation, this identity holds for $\sigma > -1$.

We have [4], p. 126,

$$(7) \quad -\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} = \int_0^{\infty} \frac{\sin \frac{xt}{2}}{e^{\pi t} - 1} dt.$$

But the sine function is enveloped by its MACLAURIN series [1], p. 26. Since $e^{\pi t} - 1 > 0$ for $t > 0$, we may use (7) to obtain the result

$$(8) \quad -\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} \propto \sum_{k=1}^{\infty} \frac{(-)^{k-1} B_k}{(2k)!} x^{2k-1} \quad (-\infty < x < \infty),$$

where the BERNOULLI numbers B_k are given by the integrals [4], p. 126,

$$(9) \quad B_k = 2^{2-2k} \int_0^{\infty} \frac{t^{2k-1} dt}{e^{\pi t} - 1}.$$

For real $s > -1$, we may use the enveloping series (8) in an integration to obtain

$$(10) \quad \int_0^{\infty} e^{-ax} x^{s-1} \left(-\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} \right) dx \propto \sum_{k=1}^{\infty} \frac{(-)^{k-1} B_k \Gamma(s+2k-1)}{(2k)! a^{s+2k-1}}.$$

From (6) we now find the enveloping series

$$(11) \quad \zeta(s, a) - \frac{1}{(s-1)a^{s-1}} - \frac{1}{2a^s} \propto \sum_{k=1}^{\infty} \frac{(-)^{k-1} B_k s(s+1) \dots (s+2k-2)}{(2k)! a^{s+2k-1}}.$$

The result (11), which holds for real $s > -1$ may be extended to real values ≤ -1 and to complex values. Let

$$(12) \quad R_n(x) = -\frac{1}{x} + \frac{1}{2} \coth \frac{x}{2} - \sum_{k=1}^n \frac{(-)^{k-1} B_k}{(2k)!} x^{2k-1}.$$

The remainder $R_n(x)$ is $O(x^{2n+1})$ at the origin. Therefore, by analytic continuation, the identity

$$(13) \quad \left\{ \int_0^{\infty} e^{-ax} x^{s-1} R_n(x) dx = \zeta(s, a) \Gamma(s) - \frac{\Gamma(s-1)}{a^{s-1}} - \frac{1}{2} \frac{\Gamma(s)}{a^s} - \sum_{k=1}^n \frac{(-)^{k-1} B_k \Gamma(s+2k-1)}{(2k)! a^{s+2k-1}} \right.$$

holds for all complex s with $\sigma > -2n-1$. But, by (8) and (12),

$$(14) \quad R_n(x) \propto \sum_{k=n+1}^{\infty} \frac{(-)^{k-1} B_k}{(2k)!} x^{2k-1} \quad (-\infty < x < \infty).$$

Using this enveloping series in (13), we find that for all real $s > -2n-1$

$$(15) \quad \zeta(s, a) = \frac{1}{(s-1)a^{s-1}} + \frac{1}{2a^s} + \sum_{k=1}^n \frac{(-)^k B_k s(s+1) \dots (s+2k-2)}{(2k)! a^{s+2k-1}} + T_n,$$

where T_n is the integral (13) divided by $\Gamma(s)$. By (14)

$$(16) \quad T_n \propto \sum_{k=n+1}^{\infty} \frac{(-)^k B_k s(s+1) \dots (s+2k-2)}{(2k)! a^{s+2k-1}}.$$

Finally suppose that s is complex, with $\sigma > -2n - 1$. In this case the identity (15) still holds, but the enveloping relation (16) fails. Using the enveloping series (14) for $R_n(x)$ in the integral (13), we find for each $N \geq n + 1$

$$(17) \quad T_n = \sum_{k=n+1}^N \frac{(-)^k B_k s(s+1) \dots (s+2k-2)}{(2k)! a^{s+2k-1}} + \psi_N \frac{B_{N+1} \Gamma(\sigma+2N+2)}{(2N+2)! \Gamma(s) a^{\sigma+2N+1}},$$

where ψ_N is a complex number with $|\psi_N| < 1$. If s is a complex number with $\sigma > -2n - 1$, the result (15) may be used where the relation (17) is substituted for the strict enveloping relation (16).

California Institute of Technology

REFERENCES

1. PÓLYA, G. and G. SZEGÖ, Aufgaben und Lehrsätze aus der Analysis, I, 26-29, Springer (1925).
2. SCHOTT, G. A. and G. N. WATSON, Asymptotic Formulae Occurring in Electron Theory, Quart. J. of Pure and Appl. Math., 47, 311-333 (1917).
3. CORPUT, J. G. VAN DER, Asymptotic Expansions, Lectures, National Bureau of Standards, Los Angeles, California, Summer Session 1951 (pre-publication copy), 178 pages. Asymptotic Expansions II, Elementary Methods, Technical Report, University of California, Berkeley, California.
4. WHITTAKER, E. T. and G. N. WATSON, A Course of Modern Analysis, American Edition, Macmillan (1948).